

Methods: From 105 patients (male gender 79%, age 61.6 ± 11.0 years), the angiographic grade and direction of collateral flow in 121 totally occluded vessels was compared to the transmural attenuation gradient of vessel distal to total occlusion (TAGdistal) derived from 64-detector row CCTA. TAG is defined as the linear gradient of luminal attenuation along coronary artery, and was validated against angiographical stenosis and flow velocity in our previous study.

Results: TAGdistal increased consistently and significantly with the degree of collateral flow, from -4.43 ± 4.02 HU/mm for Rentrop score 0 to 0.82 ± 1.08 HU/mm for Rentrop score 3 ($p < 0.0001$). TAGdistal was also significantly higher in retrograde flow compared to antegrade collateral flow (-2.44 ± 3.04 HU/mm versus 1.33 ± 2.59 HU/mm, $p < 0.0001$). The well-developed collateral vessel that have Rentrop score 2 or 3, which was found in 42.1% (51/121), could be predicted by the TAGdistal cutoff value of ≥ -1.28 HU/mm with area under receiver operating characteristic curve of 0.689, and with a sensitivity and specificity, positive and negative predictive value of 86.3%, 47.1%, and 54.3%, 82.5%, respectively.

Conclusions: As far as we know, this is the first study showing that CT can evaluate coronary collateral flow. Using TAG method, CCTA appears to be able to measure quantitatively the degree and direction of coronary collateral circulation, and predict angiographically well developed collateral vessels. These abilities of CCTA may be useful for evaluation of patients with complex coronary artery disease.

TCT-74

Impact of Pre-Procedural Coronary CT Angiography on the Procedural Success of Percutaneous Coronary Intervention for Chronic Total Occlusion: A Multicenter Study of e-CTO Investigators

Jin-Ho Choi¹, Young-Bin Song¹, Joo-Yong Hahn¹, Seung-Hyuk Choi¹, Hyeon-Cheol Gwon¹, Cheol-Woong Yoo², Hee-Yeol Kim², Seung-Woon Rha², Young Keun Ahn³, Jong Seon Park⁴, Doo-Il Kim², Seung-Ho Huh², Seung Hwan Lee⁵, Jang-Hyun Cho², Jang-Ho Bae², Donghoon Choi², In-Ho Chae², Hyo-Soo Kim⁶, Hun-Sik Park²

¹Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea, Republic of; ²e-CTO Investigators, Seoul, Korea, Republic of; ³e-CTO Investigators, Gwangju, Korea, Republic of; ⁴e-CTO Investigators, Dae Gu, Korea, Republic of; ⁵e-CTO Investigators, Wonju, Korea, Republic of; ⁶e-CTO Investigators, Seoul, South Korea

Background: Coronary CT angiography (CCTA) has been used to predict procedural success of percutaneous coronary intervention (PCI) for chronic total occlusion (CTO). However, the role of CCTA on the procedural outcome has not been reported in a large-scaled study. We investigated the impact of pre-procedural CCTA on the procedural success of CTO PCI on a patient-basis.

Methods: We retrospectively compared 2,840 patients without pre-procedural CCTA (no CCTA group) and 658 patients with CCTA (CCTA group) from e-CTO, a Korean multicenter registry comprising 26 centers. Results were further confirmed in propensity-matched subgroup (N=1,316).

Results: CCTA groups were younger (62.1 ± 10.6 vs. 63.1 ± 11.2), more were female (21.9% vs. 27.4%), more had CTO in right coronary artery (45.0% vs. 37.6%, $p < 0.05$), and more risk factors. These differences were eliminated after matching of 21 variables. The unadjusted CTO PCI success rate was lower in CCTA group compared to non CCTA group (83.4% vs. 75.2%), and it was consistent in subgroup analyses by lesion location (LAD, 78.4% vs. 85.7%; LCX, 72.5% vs. 84.3%; RCA, 73.3% vs. 80.4%, $p < 0.05$). The use of pre-procedural CCTA was related to 0.61-fold decrease of odds for procedural success (95% confidence interval (CI)=0.49-0.74, $p < 0.001$) in unadjusted model. This result was confirmed in covariate-adjusted model (OR=0.57, 95% CI=0.45 - 0.71) and in propensity-score matched model (OR=0.61, 95% CI=0.47 - 0.80, $p < 0.001$).

Conclusions: Pre-procedural CCTA did not show beneficial impact on the procedural success of CTO PCI in our multicenter registry. Careful selection or sophisticated CCTA analytic methods would be required to demonstrate the clinical role of pre-procedural CCTA before CTO PCI.

TCT-75

Lumen Enlargement of the Coronary Segments Located Distal to Chronic Total Occlusions Successfully Treated with Drug-Eluting Stents at Follow-up

Josep Gomez Lara¹, Luis Teruel¹, Silvia Homs¹, José Luis Ferreiro¹, Rafael Romaguera¹, Gerard Roura², Guillermo Sánchez-Elvira¹, Salvatore Brugaletta³, Joan Antoni Gomez-Hospital¹, Angel Cequier⁴

¹Hospital Universitari de Bellvitge, Hospitalet de Llobregat, Spain; ²Hospital Universitari Bellvitge, Hospitalet de Llobregat, Spain; ³Thorax institute, Barcelona, Spain; ⁴Hospital of Bellvitge, Barcelona, Spain

Background: Chronic total occlusions (CTO) are the final stage of coronary atherosclerosis. Coronary arteries with CTO have shown large plaque burden and negative remodeling of the occluded region and the segments located distal to the occlusion. Lumen and plaque changes located distal to successfully re-canalized CTO remain unknown at follow-up.

Methods: Ninety-one CTO successfully treated with drug-eluting stents in 86 patients underwent quantitative angiography at baseline and 12-18 months

follow-up. Thirty-one lesions were investigated with IVUS. All analyses were performed after nitroglycerin. Angiographic changes of were assessed with quantitative coronary angiography as differences in minimal, mean and maximal lumen diameter (MinLD, MeanLD and MaxLD, respectively). Vessel remodeling was assessed with IVUS as changes in lumen, plaque and vessel volume.

Results: At follow-up, MinLD increased 23.9% (from 0.88 ± 0.32 to 1.09 ± 0.35 mm; $p < 0.01$), MeanLD 16.4% (from 1.59 ± 0.44 to 1.85 ± 0.45 mm; $p < 0.01$) and MaxLD 11.7% (from 2.39 ± 0.67 to 2.67 ± 0.70 mm; $p < 0.01$). Lumen enlargement was greater in non-restenotic lesions, small lumen area at baseline and low LDL-cholesterol levels during the study period. By IVUS, lumen increased 26.9% (from 108.1 ± 89.2 to 137.3 ± 115.3 mm³; $p < 0.01$), vessel increased 12.1% (from 207.1 ± 170.2 to 232.2 ± 196.0 mm³; $p < 0.01$) and plaque tended to decrease (-3.9%, from 98.9 ± 88.7 to 94.9 ± 89.3 mm³; $p = 0.07$). Small lumen at baseline was related to greater lumen enlargement.

Conclusions: Distal segments to re-canalized CTO show a notable lumen and vessel enlargement with a trend towards of mild plaque regression. Low LDL-Cholesterol levels during the study increases lumen enlargement. Angiographic lesions distal to CTO may change and stent implantation must be discouraged.

TCT-76

Predictive Value of the J-CTO Score in Percutaneous Coronary Interventions for Chronic Total Occlusions

Giuseppe Ferrante¹, Yves Louvard¹, Talal Harb¹, Thierry Unterseeck², Thomas Hovasse¹, Marie-Claude Morice¹, Bernard Chevalier¹, Thierry Lefevre¹, Philippe Garot²

¹ICPS, Massy, France; ²ICPS, Quincy, France

Background: Introduction The J-CTO score has been shown to predict successful guidewire crossing within 30 minutes in percutaneous coronary intervention (PCI) for chronic total occlusions (CTO) in a multicentre Japanese registry. Hypothesis We assessed the hypothesis that the J-CTO score is a useful risk score for the prediction of procedural failure of PCI for CTO in a different cohort of patients.

Methods: Methods The study included all consecutive patients undergoing PCI for CTO at 3 tertiary PCI centres between January 2004 and December 2011. The J-CTO score assigns 1 point to each of the following: calcification, bending, blunt stump, occlusion length ≥ 20 mm, and previously failed lesion and classifies lesions as easy (score of 0), intermediate (score of 1), difficult (score of 2), and "very difficult" (score of ≥ 3). A multivariable mixed effect logistic regression for clustered data was used to assess the impact of J-CTO score on PCI failure. Model calibration was assessed as difference between predicted probabilities with the worst or best prognosis (PSEP). Areas under receiver-operating characteristic curve (AUC) were computed.

Results: Results A total of 1261 patients, median age 63 yrs-old (25th-75th percentile, 55-72), undergoing PCI for 1418 CTO were included. PCI failure occurred in 410 (28.9%) lesions. Failure rate significantly increased with increasing J-CTO score (13.6%, 24.7%, 37.0%, 44.8%, in the groups with J-CTO score of 0, 1, 2, ≥ 3 , respectively, $p < 0.001$). At multivariable logistic regression J-CTO score was a significant predictor of failure (odds ratio 1.68, 95% confidence interval (CI) 1.43-1.97, $p < 0.001$, for each unit increase in J-CTO score). PSEP was 0.34 and 0.33 in a model containing J-CTO score only, or containing J-CTO score in addition to clinical, procedural variables and vessel site, respectively. The AUC of a model containing J-CTO score only was significantly higher than AUC of a model containing J-CTO score in addition to clinical, procedural variables and vessel site (0.77, 95% CI 0.75-0.80, vs. 0.71, 95% CI 0.69-0.74, $p < 0.001$).

Conclusions: Conclusions The J-CTO score is an independent predictor of failure of PCI for CTO and has a good predictive accuracy as stand-

TCT-77

Initial and Mid-Term Angiographic Outcomes of Septal Channel Perforation Related to Retrograde Recanalization for Chronic Total Occlusions

Hiroyuki Tanaka¹, Kazushige Kadota¹, Seiji Habara¹, Yasushi Fuku¹, Tsuyoshi Goto¹, Kazuaki Mitsudo¹

¹Kurashiki Central Hospital, Kurashiki, Japan

Background: Septal channel perforation occurs rarely in retrograde recanalization via septal channel for chronic total occlusion (CTO) lesions. There has been little data on mid-term angiographic outcomes of septal channel perforation.

Methods: Septal channel perforation with cardiac tamponade or spreading myocardial blush was treated with coil or fat tissue to occlude, balloon dilatation at the donor artery, and administration of protamine. Persistent septal channel perforation into the ventricle and coronary sinus or of non-growing myocardial blush was followed up by no treatment. We examined the angiographic outcome of no treatment cases of septal channel perforation.

Results: Between October 2005 and December 2011, we performed the retrograde approach in 465 patients with 484 CTO lesions. Of these, the septal channel was used in 55.2% (267/484), and its in-hospital outcomes were no major adverse cardiac events and 1 cardiac tamponade. The incidence of septal channel perforation was 15.4% (41/267). In septal channel perforation cases, we used the coil (n=5), fat tissue (n=2), balloon dilatation (n=3), and protamine (n=15). Of